

LAMPIRAN

Warning # 849 in column 23. Text: in_ID
 The LOCALE subcommand of the SET command has an invalid parameter.
 It could
 not be mapped to a valid backend locale.
 CORRELATIONS
 /VARIABLES=X1 X2 X3 X4 X5 X6 X7 X8 X9 X10 X11 X12 X13 X14 X15
 TOTAL_X
 /PRINT=TWOTAIL NOSIG
 /MISSING=PAIRWISE.

Correlations

Notes		
Output Created		13-JAN-2019 19:31:32
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X1 X2 X3 X4 X5 X6 X7 X8 X9 X10 X11 X12 X13 X14 X15 TOTAL_X /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02

[DataSet0]

X10	Pearson Correlation	,314	,463*	,309	,267	,047	,443*	,470*	,328	,200
	Sig. (2-tailed)	,097	,012	,103	,161	,807	,016	,010	,083	,298
	N	29	29	29	29	29	29	29	29	29
X11	Pearson Correlation	,472**	,258	,475**	,516**	,190	,416*	,250	,430*	,230
	Sig. (2-tailed)	,010	,176	,009	,004	,325	,025	,192	,020	,230
	N	29	29	29	29	29	29	29	29	29
X12	Pearson Correlation	,602**	,416*	,577**	,248	,093	,587**	,387*	,445*	,507**
	Sig. (2-tailed)	,001	,025	,001	,195	,632	,001	,038	,016	,005
	N	29	29	29	29	29	29	29	29	29
X13	Pearson Correlation	,379*	,389*	,245	,152	-,043	,403*	,219	,250	,308
	Sig. (2-tailed)	,043	,037	,199	,431	,826	,030	,253	,192	,104
	N	29	29	29	29	29	29	29	29	29
X14	Pearson Correlation	,424*	,369*	,140	,428*	,103	,228	,281	,176	,084
	Sig. (2-tailed)	,022	,049	,468	,021	,595	,235	,140	,362	,663
	N	29	29	29	29	29	29	29	29	29
X15	Pearson Correlation	,469*	,397*	,048	,182	,163	,434*	,326	,100	,185
	Sig. (2-tailed)	,010	,033	,806	,345	,398	,019	,085	,607	,337
	N	29	29	29	29	29	29	29	29	29
TOTAL_X	Pearson Correlation	,763**	,773**	,609**	,606**	,451*	,636**	,757**	,762**	,491**
	Sig. (2-tailed)	,000	,000	,000	,000	,014	,000	,000	,000	,007
	N	29	29	29	29	29	29	29	29	29

Correlations

		X10	X11	X12	X13	X14	X15	TOTAL_X
X1	Pearson Correlation	,314	,472**	,602**	,379*	,424*	,469*	,763**
	Sig. (2-tailed)	,097	,010	,001	,043	,022	,010	,000
	N	29	29	29	29	29	29	29
X2	Pearson Correlation	,463*	,258	,416*	,389*	,369*	,397*	,773**
	Sig. (2-tailed)	,012	,176	,025	,037	,049	,033	,000
	N	29	29	29	29	29	29	29

X3	Pearson Correlation	,309	,475**	,577**	,245	,140	,048	,609**
	Sig. (2-tailed)	,103	,009	,001	,199	,468	,806	,000
	N	29	29	29	29	29	29	29
X4	Pearson Correlation	,267	,516**	,248	,152	,428*	,182	,606**
	Sig. (2-tailed)	,161	,004	,195	,431	,021	,345	,000
	N	29	29	29	29	29	29	29
X5	Pearson Correlation	,047	,190	,093	-,043	,103	,163	,451*
	Sig. (2-tailed)	,807	,325	,632	,826	,595	,398	,014
	N	29	29	29	29	29	29	29
X6	Pearson Correlation	,443*	,416*	,587**	,403*	,228	,434*	,636**
	Sig. (2-tailed)	,016	,025	,001	,030	,235	,019	,000
	N	29	29	29	29	29	29	29
X7	Pearson Correlation	,470*	,250	,387*	,219	,281	,326	,757**
	Sig. (2-tailed)	,010	,192	,038	,253	,140	,085	,000
	N	29	29	29	29	29	29	29
X8	Pearson Correlation	,328	,430*	,445*	,250	,176	,100	,762**
	Sig. (2-tailed)	,083	,020	,016	,192	,362	,607	,000
	N	29	29	29	29	29	29	29
X9	Pearson Correlation	,200	,230	,507**	,308	,084	,185	,491**
	Sig. (2-tailed)	,298	,230	,005	,104	,663	,337	,007
	N	29	29	29	29	29	29	29
X10	Pearson Correlation	1	,453*	,500**	,693**	,460*	,423*	,659**
	Sig. (2-tailed)		,014	,006	,000	,012	,022	,000
	N	29	29	29	29	29	29	29
X11	Pearson Correlation	,453*	1	,710**	,670**	,216	,022	,677**
	Sig. (2-tailed)	,014		,000	,000	,260	,909	,000
	N	29	29	29	29	29	29	29
X12	Pearson Correlation	,500**	,710**	1	,710**	,231	,269	,762**
	Sig. (2-tailed)	,006	,000		,000	,227	,159	,000
	N	29	29	29	29	29	29	29
X13	Pearson Correlation	,693**	,670**	,710**	1	,334	,239	,628**
	Sig. (2-tailed)	,000	,000	,000		,077	,212	,000
	N	29	29	29	29	29	29	29
X14	Pearson Correlation	,460*	,216	,231	,334	1	,748**	,525**
	Sig. (2-tailed)	,012	,260	,227	,077		,000	,003
	N	29	29	29	29	29	29	29
X15	Pearson Correlation	,423*	,022	,269	,239	,748**	1	,490**
	Sig. (2-tailed)							
	N							

	Sig. (2-tailed)	,022	,909	,159	,212	,000		,007
	N	29	29	29	29	29	29	29
TOTAL_X	Pearson Correlation	,659**	,677**	,762**	,628**	,525**	,490**	1
	Sig. (2-tailed)	,000	,000	,000	,000	,003	,007	
	N	29	29	29	29	29	29	29

*. Correlation is significant at the 0.05 level (2-tailed).

** . Correlation is significant at the 0.01 level (2-tailed).

RELIABILITY

```

/VARIABLES=X1 X2 X3 X4 X5 X6 X7 X8 X9 X10 X11 X12 X13 X14 X15
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA.

```

Reliability

Notes

Output Created		13-JAN-2019 19:32:09
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
Missing Value Handling	Matrix Input	
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.
Syntax		RELIABILITY /VARIABLES=X1 X2 X3 X4 X5 X6 X7 X8 X9 X10 X11 X12 X13 X14 X15 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,00

Scale: ALL VARIABLES

Case Processing Summary

		N	%
Cases	Valid	29	100,0
	Excluded ^a	0	,0
	Total	29	100,0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
,896	15

```
NEW FILE.
DATASET NAME DataSet1 WINDOW=FRONT.
CORRELATIONS
  /VARIABLES=Y16 Y17 Y18 Y19 Y20 Y21 Y22 Y23 Y24 Y25 Y26 Y27 Y28
Y29 Y30 TOTAL_Y
  /PRINT=TWOTAIL NOSIG
  /MISSING=PAIRWISE.
```

Correlations

Notes

Output Created		13-JAN-2019 19:37:33
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.

Correlations

		Y25	Y26	Y27	Y28	Y29	Y30	TOTAL_ Y
Y16	Pearson Correlation	,355	,370 ⁺	,444 [*]	,267	,175	,397 ⁺	,645 ^{**}
	Sig. (2-tailed)	,059	,048	,016	,161	,364	,033	,000
	N	29	29	29	29	29	29	29
Y17	Pearson Correlation	,135	,014	,146	,355	,290	,276	,442 ⁺
	Sig. (2-tailed)	,485	,943	,451	,059	,127	,147	,016
	N	29	29	29	29	29	29	29
Y18	Pearson Correlation	,051	,460 ⁺	,313	,193	,316	,029	,653 ^{**}
	Sig. (2-tailed)	,792	,012	,098	,317	,095	,882	,000
	N	29	29	29	29	29	29	29
Y19	Pearson Correlation	,111	,517 ^{**}	,516 ^{**}	,060	,073	,022	,529 ^{**}
	Sig. (2-tailed)	,566	,004	,004	,756	,707	,909	,003
	N	29	29	29	29	29	29	29
Y20	Pearson Correlation	,234	,322	,389 ⁺	,098	,152	,098	,668 ^{**}
	Sig. (2-tailed)	,221	,088	,037	,613	,431	,611	,000
	N	29	29	29	29	29	29	29
Y21	Pearson Correlation	-,077	,025	,139	,243	,012	,112	,426 ⁺
	Sig. (2-tailed)	,690	,898	,471	,205	,952	,562	,021
	N	29	29	29	29	29	29	29
Y22	Pearson Correlation	,102	,326	,222	,247	,305	,238	,587 ^{**}
	Sig. (2-tailed)	,600	,085	,248	,196	,107	,213	,001
	N	29	29	29	29	29	29	29
Y23	Pearson Correlation	,292	,326	,222	,318	,305	,238	,596 ^{**}
	Sig. (2-tailed)	,125	,085	,248	,093	,107	,213	,001
	N	29	29	29	29	29	29	29
Y24	Pearson Correlation	,422 ⁺	,419 ⁺	,457 ⁺	,203	,258	,009	,636 ^{**}
	Sig. (2-tailed)	,023	,024	,013	,291	,177	,962	,000
	N	29	29	29	29	29	29	29
Y25	Pearson Correlation	1	,498 ^{**}	,339	,336	,398 ⁺	,240	,493 ^{**}
	Sig. (2-tailed)		,006	,072	,074	,032	,209	,007
	N	29	29	29	29	29	29	29
Y26	Pearson Correlation	,498 ^{**}	1	,680 ^{**}	,092	,335	,203	,625 ^{**}
	Sig. (2-tailed)	,006		,000	,634	,075	,292	,000
	N	29	29	29	29	29	29	29
Y27	Pearson Correlation	,339	,680 ^{**}	1	,149	,228	,453 ⁺	,639 ^{**}

	Sig. (2-tailed)	,072	,000		,439	,234	,014	,000
	N	29	29	29	29	29	29	29
Y28	Pearson Correlation	,336	,092	,149	1	,706**	,248	,540**
	Sig. (2-tailed)	,074	,634	,439		,000	,194	,002
	N	29	29	29	29	29	29	29
Y29	Pearson Correlation	,398*	,335	,228	,706**	1	,461*	,592**
	Sig. (2-tailed)	,032	,075	,234	,000		,012	,001
	N	29	29	29	29	29	29	29
Y30	Pearson Correlation	,240	,203	,453*	,248	,461*	1	,442*
	Sig. (2-tailed)	,209	,292	,014	,194	,012		,016
	N	29	29	29	29	29	29	29
TOTAL_Y	Pearson Correlation	,493**	,625**	,639**	,540**	,592**	,442*	1
	Sig. (2-tailed)	,007	,000	,000	,002	,001	,016	
	N	29	29	29	29	29	29	29

*. Correlation is significant at the 0.05 level (2-tailed).

**. Correlation is significant at the 0.01 level (2-tailed).

RELIABILITY

```

/VARIABLES=Y16 Y17 Y18 Y19 Y20 Y21 Y22 Y23 Y24 Y25 Y26 Y27 Y28
Y29 Y30
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.

```

Reliability

Notes

Output Created		13-JAN-2019 19:38:03
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Matrix Input	
	Definition of Missing	User-defined missing values are treated as missing.

	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.
Syntax		RELIABILITY /VARIABLES=Y16 Y17 Y18 Y19 Y20 Y21 Y22 Y23 Y24 Y25 Y26 Y27 Y28 Y29 Y30 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01

Scale: ALL VARIABLES

Case Processing Summary

		N	%
Cases	Valid	29	100,0
	Excluded ^a	0	,0
	Total	29	100,0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
,843	15

Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
Y16	57,97	29,034	,564	,827
Y17	57,83	31,576	,368	,839
Y18	58,41	27,466	,539	,830
Y19	58,38	29,458	,412	,838
Y20	58,21	28,241	,579	,826

Y21	58,34	31,448	,342	,840
Y22	58,07	29,995	,508	,831
Y23	58,07	29,924	,519	,831
Y24	58,14	28,623	,542	,829
Y25	58,03	30,892	,410	,837
Y26	58,17	31,505	,586	,833
Y27	58,17	30,362	,582	,829
Y28	58,24	29,547	,432	,836
Y29	58,17	29,362	,500	,831
Y30	58,14	30,980	,344	,840

```

NEW FILE.
DATASET NAME DataSet2 WINDOW=FRONT.
REGRESSION
  /MISSING LISTWISE
  /STATISTICS COEFF OUTS R ANOVA
  /CRITERIA=PIN(.05) POUT(.10)
  /NOORIGIN
  /DEPENDENT Y
  /METHOD=ENTER X
  /RESIDUALS NORMPROB(ZRESID) .

```

Regression

Notes

Output Created		13-JAN-2019 19:41:02
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any variable used.

Syntax	REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT Y /METHOD=ENTER X /RESIDUALS NORMPROB(ZRESID).	
Resources	Processor Time	00:00:01,59
	Elapsed Time	00:00:00,94
	Memory Required	1356 bytes
	Additional Memory Required for Residual Plots	312 bytes

[DataSet2]

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	Sistem Informasi Akuntansi ^b		Enter

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

b. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,647 ^a	,419	,397	4,523

a. Predictors: (Constant), Sistem Informasi Akuntansi

b. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	397,761	1	397,761	19,440	,000 ^b
	Residual	552,446	27	20,461		
	Total	950,207	28			

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

b. Predictors: (Constant), Sistem Informasi Akuntansi

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t
		B	Std. Error	Beta	
1	(Constant)	29,214	7,553		3,868
	Sistem Informasi Akuntansi	,529	,120	,647	4,409

Coefficients^a

Model		Sig.
1	(Constant)	,001
	Sistem Informasi Akuntansi	,000


a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

Residuals Statistics^a

	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	50,92	67,33	62,31	3,769	29
Residual	-8,331	12,081	,000	4,442	29
Std. Predicted Value	-3,022	1,332	,000	1,000	29
Std. Residual	-1,842	2,671	,000	,982	29

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

Charts



```

REGRESSION
  /MISSING LISTWISE
  /STATISTICS COEFF OUTS R ANOVA
  /CRITERIA=PIN(.05) POUT(.10)
  /NOORIGIN
  /DEPENDENT Y
  /METHOD=ENTER X
  /RESIDUALS NORMPROB(ZRESID)
  /SAVE RESID.
  
```

Regression

Notes

Output Created	13-JAN-2019 19:41:44	
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	
	File	29

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any variable used.
Syntax		REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT Y /METHOD=ENTER X /RESIDUALS NORMPROB(ZRESID) /SAVE RESID.
Resources	Processor Time	00:00:00,28
	Elapsed Time	00:00:00,31
	Memory Required	1356 bytes
	Additional Memory Required for Residual Plots	312 bytes
Variables Created or Modified	RES_1	Unstandardized Residual

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	Sistem Informasi Akuntansi	.	Enter

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

b. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,647 ^a	,419	,397	4,523

a. Predictors: (Constant), Sistem Informasi Akuntansi

b. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	397,761	1	397,761	19,440	,000 ^b
	Residual	552,446	27	20,461		
	Total	950,207	28			

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

b. Predictors: (Constant), Sistem Informasi Akuntansi

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t
		B	Std. Error	Beta	
1	(Constant)	29,214	7,553		3,868
	Sistem Informasi Akuntansi	,529	,120	,647	4,409

Coefficients^a

Model		Sig.
1	(Constant)	,001
	Sistem Informasi Akuntansi	,000


a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

Residuals Statistics^a

	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	50,92	67,33	62,31	3,769	29
Residual	-8,331	12,081	,000	4,442	29
Std. Predicted Value	-3,022	1,332	,000	1,000	29
Std. Residual	-1,842	2,671	,000	,982	29

a. Dependent Variable: Pengendalian Intern Pembelian Bahan Baku

Charts



```

NPAR TESTS
  /K-S (NORMAL)=RES_1
  /MISSING ANALYSIS.
  
```

NPar Tests

Notes

Output Created		13-JAN-2019 19:42:30
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each test are based on all cases with valid data for the variable(s) used in that test.
Syntax		NPAR TESTS /K-S(NORMAL)=RES_1 /MISSING ANALYSIS.
Resources	Processor Time	00:00:00,00

Elapsed Time	00:00:00,01
Number of Cases Allowed ^a	196608

a. Based on availability of workspace memory.

One-Sample Kolmogorov-Smirnov Test

		Unstandardized Residual
N		29
Normal Parameters ^{a,b}	Mean	,0000000
	Std. Deviation	4,44186966
Most Extreme Differences	Absolute	,111
	Positive	,111
	Negative	-,064
Test Statistic		,111
Asymp. Sig. (2-tailed)		,200 ^{c,d}

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

d. This is a lower bound of the true significance.

```

CORRELATIONS
/VARIABLES=X Y
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

Correlations

Notes

Output Created		13-JAN-2019 19:44:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data	29
	File	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.

Cases Used		Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X Y /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,01

Correlations

		Sistem Informasi Akuntansi	Pengendalian Intern Pembelian Bahan Baku
Sistem Informasi Akuntansi	Pearson Correlation	1	,647**
	Sig. (2-tailed)		,000
	N	29	29
Pengendalian Intern Pembelian Bahan Baku	Pearson Correlation	,647**	1
	Sig. (2-tailed)	,000	
	N	29	29

** . Correlation is significant at the 0.01 level (2-tailed).

```
DATASET ACTIVATE DataSet0.
```

```
SAVE OUTFILE='C:\Users\ACER\Documents\SPSS\TABEL X.sav'  
/COMPRESSED.
```

```
DATASET ACTIVATE DataSet1.
```

```
SAVE OUTFILE='C:\Users\ACER\Documents\SPSS\TABEL Y.sav'  
/COMPRESSED.
```

```
DATASET ACTIVATE DataSet2.
```

```
SAVE OUTFILE='C:\Users\ACER\Documents\SPSS\TABEL XY.sav'  
/COMPRESSED.
```

```
DATASET ACTIVATE DataSet0.
```

```
DATASET CLOSE DataSet2.
```

```
DATASET ACTIVATE DataSet1.
```

```
DATASET CLOSE DataSet0.
```